




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 15/00, C07K 7/10 A61K 39/12, 39/42, G01N 33/566 G01N 33/577	A1	(11) International Publication Number: WO 92/13946 (43) International Publication Date: 20 August 1992 (20.08.92)
(21) International Application Number: PCT/US92/00823 (22) International Filing Date: 3 February 1992 (03.02.92) (30) Priority data: 653,091 8 February 1991 (08.02.91) US (71) Applicant: GENELABS INCORPORATED [US/US]; 505 Penobscot Drive, Redwood City, CA 94063 (US). (72) Inventors: REYES, Gregory, R. ; 780 Palo Alto Avenue, Palo Alto, CA 94301 (US). HADLOCK, Kenneth, G. ; 22815 Vermont Street, #501, Hayward, CA 94541 (US). (74) Agent: DEHLINGER, Peter, J.; P.O. Box 60850, Palo Alto, CA 94306-1546 (US).	(81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, KR, LU (European patent), MC (European patent), NL (European patent), SE (European patent). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	

(54) Title: HTLV-I AND HTLV-II PEPTIDE ANTIGENS AND METHODS**(57) Abstract**

Novel HTLV-I and HTLV-II peptide antigens are disclosed for use in diagnostics assays for screening and confirming HTLV-I and HTLV-II antisera. The peptides are derived from analogous regions of HTLV-I and HTLV-II gp 46 envelope proteins, and are differentiated by their immunoreactivity with an HTLV-II specific monoclonal antibody and by HTLV-I and HTLV-II antisera. The peptides are also useful in vaccine compositions.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	MI	Mali
AU	Australia	FR	France	MN	Mongolia
BB	Barbados	GA	Gabon	MR	Mauritania
BE	Belgium	GB	United Kingdom	MW	Malawi
BF	Burkina Faso	GN	Guinea	NL	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	RO	Romania
CA	Canada	IT	Italy	RU	Russian Federation
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark	MG	Madagascar		
ES	Spain				

5

HTLV-I AND HTLV-II PEPTIDE ANTIGENS AND METHODS

This application is a continuation-in-part of co-pending U.S. patent application for "HTLV-I Peptide
10 Antigen and Methods", Serial No. 366,313, filed June 13, 1989, which in turn is a continuation of U. S. patent application for "HTLV-I Peptide Antigen and Methods," Serial No. 948,270, filed December 31, 1986, now abandoned.

15

1. Field of the Invention

The present invention relates to an HTLV-I specific antigen, and to methods of preparing and using the antigen.

20

2. References

Cwirla, S.E., et al., Proc Nat Acad. Sci, USA, 87:6378 (1990).

Huynh, T.V., et al., in "DNA Cloning, Volume 1,"
25 ed. D.M.Glover, Washington, D.C.: IRL Press, 1985 (Chapter 2).

Laemmli, U.K., Nature, 227:680 (1970).

Lipka, J.J., et al., J Infect Dis, 162:353 (1990).

Lipka, J.J., et al., Proceedings of the 43 Meeting
30 (1990).

Maniatis, T., et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory (1982).

Matsushita, S., et al., Proc Natl Acad Sci (USA),
5 83:2672 (1986).

Miyoshi, I, et al., Nature, 294:770 (1981).

Poiesz, B.J., et al., Proc Natl Acad Sci (USA),
77:7415 (1980).

Popovic, M., et al., Science, 29:856 (1983).

10 Samuel, K.P., et al., Science, 226:1094 (1984).

Samuel, K.P., Gene Anal Tech, 2:60 (1985)

Scott, J.T. et al., Science, 249:386 (1990).

Seiki, M., et al., Proc Natl Acad Sci (USA),
80:3618 (1983).

15 Shimotokno, K., et al, Proc Nat Acad Sci, USA,
82:3101 (1985).

3. Background of the Invention

The human T-cell leukemia viruses (HTLV) represent
20 a family of T-cell retroviruses with three known mem-
bers. HTLV type I (HTLV-I) has transforming activity
in vitro and is etiologically linked to adult T-cell
leukemia, which is known to be endemic in several
parts of the world. HTLV-II is another retrovirus
25 having transforming capacity *in vitro*, and has been
isolated from a patient with a T-cell variant of hairy
cell leukemia. HTLV-III, which has also been called
lymphadenopathy-associated virus and is now known as
the human immunodeficiency virus (HIV), is lytic for
30 certain kinds of T cells and has been linked to the

etiology of acquired immunodeficiency syndrome (AIDS). Unlike the HTLV-I and -II viruses, HTLV-III is not known to have *in vitro* transforming activity.

The diagnosis of HTLV-I infection is usually based on serum antibody response to HTLV-I peptide antigens. This usually involves an initial screening assay to identify HTLV-I antibodies, based on an enzyme immunoassay (EIA) with HTLV-I virion peptides. The assays presently used for blood screening detect about 0.5 to 10 0.05% HTLV-I and HTLV-II positives; of these, about 4 out of 5 are false positives. Therefore, positive sera must be further tested in a confirmatory assay, using Western blot or radioimmunoprecipitation assays which detect antibody reaction to specific HTLV-I peptide antigens. 15

Current blood testing procedures require confirmation tests based on immunoreaction with HTLV-I p24 gag protein and at least one of the envelop proteins gp46, gp21, or gp68. When the test antigens are prepared from virion proteins, only gp46 gives a high rate of antibody reaction with true HTLV-I seropositives. Even then, the reaction with gp46 may be detected only by additional antigen testing with a more sensitive radioimmunoprecipitation assay. The above screening 20 and confirmation testing identifies HTLV-I and HTLV-II positives, but does not distinguish between the two HTLV viruses.

It would therefore be desirable to provide an improved method for detecting HTLV-I positive sera. 30 In particular, the improved test should be capable of

detecting all HTLV-I and HTLV-II positive sera, with a minimum number of false positives, and also be able to distinguish HTLV-I from HTLV-II positive sera.

5 4. Summary of the Invention

It is therefore one object of the invention to provide an improved method and kit for detecting HTLV-I and HTLV-II positive human sera.

Another object of the invention is to provide such
10 method and kit capable of distinguishing HTLV-I and HTLV-II positive sera.

In the above-cited patent application for "HTLV-I Peptide Antigen and Assay," there is disclosed an HTLV-I peptide composed of a region of the HTLV-I gp46
15 envelop protein which is immunoreactive with the .5 α monoclonal antibody (Mab) produced by ATCC cell line HB8755 (Matsushita). The region is contained in a 42 amino acid sequence overlap of three gp46 peptide antigens, designated MTA-1, MTA-4, and MTA-5. The 42
20 amino acid sequence overlap region contains the sequence Ser-Leu-Leu-Val-Asp-Ala-Pro-Gly-Tyr-Asp-Pro-Ile-Trp-Phe-Leu-Asn-Thr-Glu-Pro-Ser-Gln-Leu-Pro-Pro-Thr-Ala-Pro-Pro-Leu-Leu-Pro-His-Ser-Asn-Leu-Asp-His-Ile-Leu-Glu-Pro-Ser, and may include the additional
25 residues Ile-Pro-Trp-Lys-Ser-Lys at the C-terminal Ser residue of the 42 amino acid sequence. A common amino acid sequence in recombinant and synthetic peptides which is immunoreactive with the .5 α Mab is the sequence Thr-Ala-Pro-Pro-Leu-Leu-Pro-His-Ser-Asn-Leu-
30 Asp-His-Ile-Leu-Glu-Pro-Ser.

In another aspect, the invention includes a kit for detecting the presence of HTLV-I infection in human serum. The kit includes a solid support on which the gp46 peptide antigen is carried, and a
5 reporter system for detecting the presence of human antibodies bound to the peptide antigen.

In one embodiment, the kit is in an EIA format for screening human sera for HTLV-I antibodies. In another embodiment, the peptide antigen is immobilized on
10 a strip, along with one or more confirmatory HTLV-I antigens, in a Western blot format for confirming HTLV-I serum antibodies.

In still another embodiment, the kit includes an HTLV-II specific antigen, defined below, capable of
15 reacting specifically with antibodies from HTLV-II positive sera. The kit allows for specific detection of HTLV-I and HTLV-II positive sera.

Also included in the invention is a method of detecting HTLV-I positive human sera. In this method,
20 test sera is reacted with a peptide antigen which is immunoreactive with anti-HTLV-I monoclonal antibody (Mab) derived from ATCC cell line HB8755, designated .5 α Mab. The presence of anti-HTLV-I antibodies bound to the antigen is detected by a suitable reporter-
25 labeled anti-human antibody.

The .5 α Mab-reactive peptide may be produced by a random-sequence selection method in which a mixture of random-sequence polynucleotides, preferably encoding
5-10 amino acid residues, is introduced into a suit-
30 able expression vector, to form a library of random-

sequence vectors. The expression products of the library vectors are screened for the presence of an amino acid sequence which is immunoreactive with the .5 α Mab. The library clone which expresses such an
5 immunoreactive amino acid sequence is then isolated and used for producing the polypeptide encoded by the inserted sequence.

Also disclosed herein is an HTLV-II peptide antigen comprising less than about 50 amino acids derived
10 from HTLV-II envelope protein gp46, and including the immunogenic region formed by the amino acid sequence Met-Thr-Leu-Leu-Val-Asp-Ala-Pro-Gly-Tyr-Asp-Pro-Leu-Trp-Phe-Ile-Thr-Ser-Glu-Pro-Thr-Gln-Pro-Pro-Pro-Thr-Ser-Pro-Pro-Leu-Val-His-Asp-Ser-Asp-Leu-Glu-His-Val-
15 Leu-Thr-Pro-Ser-Thr-Ser-Trp-Thr-Thr-Lys. A common amino acid sequence in recombinant and synthetic peptides which is immunoreactive with HTLV-II antisera has the sequence Ser-Pro-Pro-Leu-Val-His-Asp-Ser-Asp-Leu-Glu-His-Val-Leu-Thr-Pro-Ser or the same sequence
20 extended at the Ser C-terminus by the amino acid sequence Thr-Ser-Trp-Thr-Thr-Lys.

The peptide antigen is used in a test kit for detecting the presence of HTLV-II infection in a human serum. The kit includes a solid support which carries
25 the peptide antigen, and a reporter system for detecting the presence of human antibodies bound to the peptide antigen.

These and other objects and features of the present invention will become more fully apparent when

the following detailed description of the invention is read in conjunction with the accompanying drawings.

Brief Description of the Drawings

5 Figure 1A shows, in the upper line, a portion of the HTLV-I genome containing the gp46 envelop protein coding sequence, and in the lower line, a portion of the gp46 coding region containing the sequences which encode overlapping HTLV-I peptide antigens formed in
10 accordance with the invention, and designated MTA-4, MTA-1, and MTA-5 in Figure 1B;

Figure 2 shows the HTLV-I coding sequences and corresponding amino acid sequences for a portion of the HTLV-I envelop protein;

15 Figure 3 shows amino acid sequences of homologous regions of HTLV-I and HTLV-II gp46 in the region of the peptide antigen of the invention, and peptide sequences of several HTLV-I gp46 peptide antigens (upper part of figure) and HTLV-II peptide antigens
20 (lower part of figure) in accordance with the invention;

Figures 4A and 4B show antigenicity plots for the MTA-1 peptide and corresponding HTLV-II gp46 peptide;

25 Figure 5 illustrates recombinant methods for producing and selecting random-sequence peptides, in accordance with the invention;

Figure 6 shows the HTLV-II coding sequence, and corresponding amino acid sequence in the region of the gp46 envelop protein from which HTLV-II peptides of
30 the invention are derived; and

Figure 7 shows modified Western blots containing HTLV-I viral lysate and recombinant proteins p21E and MTA-4, where lanes A-F and G-R are HTLV-I and HTLV-II antisera, respectively.

5

Detailed Description of the Invention

I. Preparing HTLV-I Peptide Antigens

This section describes the preparation of HTLV-I peptide antigens which are immunoreactive with anti-HTLV-I antibodies found in individuals with HTLV-I-related T-cell leukemia. The antigens are prepared using random HTLV-I gene sequences 100-300 base pairs in length cloned in a suitable expression vector, then selected with antibody for expression of immunoreactive peptides.

A. HTLV-I Genomic Libraries

Genomic libraries of HTLV-I are prepared conventionally from cellular DNA containing an HTLV-I proviral genome. Duplex DNA may be prepared from HTLV-I infected cells, including T-cells isolated from patients known to be infected with HTLV-I virus, or known cell lines, such as HUT 102-B2 (Poiesz), MT-2 (Miyoshi), and MJ-tumor (Popovic) cells, all of which have been shown to produce HTLV-I virus. The viral genome is integrated into host DNA in these cells. Methods for preparing cell lines containing the HTLV-I genome are detailed in the above references.

The total host genomic DNA from the above cell line is partially digested with a frequent cutter, such as HaeIII or AluI, under conditions which produce

partial digest fragments in the 15-20 kbase size range, and the digested material is fractionated, for example, by sucrose gradient centrifugation, to isolate the 15-20 kbase fragments. The fragments are then cloned into a suitable cloning vector, preferably a phage cloning vector which can efficiently incorporate a 15-20 kbase insert. In a preferred method, the isolated fragments are treated with EcoRI methylase, and EcoRI linkers are ligated to their ends under standard conditions (Maniatis), and then cloned into a phage vector, such as λ Charon 4a, having a unique EcoRI insertion site.

The cloned genomic fragments are screened with a probe which is complementary to a selected sequence of the full-copy HTLV-I genome. HTLV-I sequences are known (Seiki), as are methods for producing radiolabeled synthetic oligonucleotide probes for selected sequences. In addition, synthetic oligonucleotides of specified sequences can be made by commercial services, such as provided by Synthetic Genetics, Inc. (San Diego, CA). Using such an oligonucleotide probe, molecular clones containing HTLV-I sequences are isolated from the library by standard hybridization procedures (Maniatis, p. 322). The clones can first be analyzed by restriction site analysis, to confirm that the full viral genomic sequence is present, as indicated by the presence of direct long terminal repeats which flank the integrated viral genome. The identified molecular clone is digested with a suitable endonuclease to release the full-copy viral genome. A preferred endonuclease for this purpose is SacI, which cuts the viral genome in the long terminal repeats (LTR)

10

at either end of the viral coding sequences, but does not produce internal cleavage. If the clonal HTLV-I genome is a variant with a third SacI site, an appropriate restriction enzyme will be chosen to isolate the full-length genome. The purified full-copy sequence is about a 9.5 kilobase fragment. Alternatively, a fragment of the genome representing the env gene sequences alone may be purified for production of the expression library.

Alternatively, cloning vectors containing full-copy HTLV-I duplex DNA have been reported (Seiki) and may be obtained directly from the investigators, as indicated in Example 1.

To produce the desired HTLV-I genomic library, the full-copy HTLV-I insert is excised from the above cloning vector, such as by complete digestion with SacI, and isolated as a 9.5 kilobase fragment, as described in Example 1. The isolated full-copy fragment is digested to produce DNA fragments, and preferably random fragments with sizes predominantly between about 100-300 base pairs. Example 1 describes the preparation of such fragments by DNAase digestion. Because it is desired to obtain peptide antigens of between about 30-100 amino acids, the digest fragments are preferably size fractionated, for example by gel electrophoresis, to select those in the approximately 100-300 base pair size range.

The genomic digest fragments are inserted into a suitable cloning vector, preferably an expression vector which permits expression of the coded-for peptide in a suitable host. One preferred expression vector is λ gt11, which contains a unique EcoRI insertion site 53 base pairs upstream of the translation termination codon of the β -galactosidase gene. Thus, the inserted sequence

will be expressed as a β -galactosidase gene. Thus, the inserted sequence will be expressed as a β -galactosidase fusion protein which contains most of the N-terminal portion of the β -galactosidase gene, the heterologous peptide, and at least a portion of the C-terminal region of the β -galactosidase gene. This vector also produces a temperature-sensitive repressor (cI857) which causes viral lysogeny at permissive temperatures, e.g., 32°C, and leads to viral lysis at elevated temperatures, e.g., 42°C. Advantages of this vector include: (1) highly efficient recombinant generation, (2) ability to select lysogenized host cells on the basis of host-cell growth at permissive, but not non-permissive temperatures, and (3) high levels of recombinant fusion protein production. Further, since phage containing a heterologous insert produce an inactive β -galactosidase enzyme, phage with inserts can be readily identified by a β -galactosidase colored-substrate reaction.

The digest fragments inserted into the expression vector may be modified, if needed, to contain selected restriction-site linkers, such as EcoRI linkers, according to conventional procedures. Example I illustrates methods for cloning the digest fragments into λ gt11, which includes the steps of blunt-ending the fragments, adding EcoRI linkers and ligating the fragments with EcoRI cut λ gt11. The resulting viral genome library may be checked to confirm that a relatively large (representative) library has been produced. This can be done, in the case of the λ gt11 vector, by infecting a suitable bacterial host, plating the bacteria, and examining the plaques for loss of β -galactosidase activity.

12

Using the procedures described in Example 1, about 60% of the plaques showed loss of enzyme activity, when compared to the level of background phage showing loss of enzyme activity, as seen in Example 1.

5 B. Peptide Antigen Expression

The genomic library formed above is screened for production of peptide antigen (expressed as a fusion protein) which is immunoreactive with the human anti-HTLV-I antibody of interest. One antibody of particular interest for diagnosing HTLV-I infection is the .5 α monoclonal antibody (Mab) which, as noted above, is has the same specificity as antibodies present in patients with T-cell leukemia related to HTLV-I infection. The antibody is produced by the EBV-transformed B-lymphocyte cell line having ATCC Deposit No. HC8755 (See Example 2).

In a preferred screening method, host cells infected with phage library vectors are plated, as above, and the plate is blotted with a nitrocellulose filter, to transfer recombinant antigens produced by the cells onto the filter. The filter is then reacted with the anti-HTLV-I antibody, washed to remove unbound antibody, and reacted with reporter-labeled, anti-human antibody, which becomes bound to the filter, in sandwich fashion, through the anti-HTLV-I antibody.

Typically, phage plaques which are identified by virtue of their production of recombinant antigen of interest are re-examined at a relatively low density, for production of antibody-reactive fusion protein. The screening procedures described in Example 2 are illustrative. Several recombinant phage clones which produced immunoreactive recombinant antigen were identified in the procedure.

The one or more library vectors identified as above are preferably analyzed by nucleic acid sequencing, to determine the positions of the peptide-coding regions within the HTLV-I genome. Methods for excising the

5 heterologous insert (including adjacent coding sequences of the fusion protein, if desired) from the selected library vectors, and for purifying and sequencing the excised fragments generally follow known procedures, as outlined in Example 3. The coding sequences of three

10 peptides which were found to be immunoreactive with the .5 α Mab are shown in the drawing. The three heterologous sequences were matched with the known sequence of HTLV-I (Seiki). As discussed more fully in Example 3, all of the sequences fall within base pairs 5565 and 5895 of the

15 HTLV-I genome, within the gene coding for the HTLV-I envelope protein gp46 (drawings, part A), and have an overlapping coding sequence (defined by the two arrows in the drawing) between base pairs 5664 and 5790 (drawing, part B). As seen in the drawing, part C, the overlapping

20 sequence codes for a 42-amino-acid peptide antigen having the following amino acid sequence:

Ser-Leu-Leu-Val-Asp-Ala-Pro-Gly-Tyr-Asp-Pro-Ile-Trp-Phe-Leu-Asn-Thr-Glu-Pro-Ser-Gln-Leu-Pro-Pro-Thr-Ala-Pro-Pro-Leu-Leu-Pro-His-Ser-Asn-Leu-Asp-His-Ile-Leu-Glu-

25 Pro-Ser. Screening studies conducted in support of the invention indicate that the MTA-1 peptide picks up the highest percentage of HTLV-I positive sera, particularly among subjects of Japanese ancestry. As seen in Figure 3, the MTA-1 peptide includes the additional Ile-Pro-Trp-

30 Lys-Ser-Lys residues at the Ser C terminus of the above sequence. In a preferred embodiment of the invention, the HTLV-I specific peptide contains the immunogenic

14

region of the C-terminal 48 amino acid MTA-1 sequence which is immunoreactive with the .5 α Mab.

More generally, the HTLV-I peptides of the invention include the immunogenic region of the above amino acid sequence which is immunoreactive with the .5 α Mab. As defined herein, the specified sequence includes minor, neutral amino substitutions which do not appreciably decrease the immunoreactivity of the peptide antigen for the .5 α Mab. Such amino substitutions may be selected on the basis of similarities in hydrophobicity, size, charge, hydrogen bonding ability, and effect on secondary structure according to known amino acid substitution principles.

The selected clones are used for scale-up production, for purposes of recombinant protein purification. Scale-up production is carried out using one of a variety of reported methods for (a) lysogenizing a suitable host, such as E. coli, with a selected λ gt11 recombinant, (b) culturing the transduced cells under conditions that yield high levels of the heterologous peptide, and (c) purifying the recombinant antigen from the lysed cells.

In one preferred method involving the above λ gt11 cloning vector, a high-producer E. coli host, BNN103, is infected with the selected library phage, and replica-plated on two plates. One of the plates is grown at 32°C, at which viral lysogeny can occur, and the other at 42°C, at which the infecting phage is in a lytic stage and therefore prevents cell growth. Cells which grow at the lower, but not the higher temperature, are therefore assumed to be successfully lysogenized.

The lysogenized host cells are then grown under liquid culture conditions which favor high production of the fused protein containing the viral insert, and lysed by rapid freezing to release the desired fusion protein.

5 These methods are detailed in Example 4.

HTLV-I coding sequences from the λ gt11 clone expressing the peptide antigen MTA-1 have been prepared by PCR amplification, as described in Section II below, and cloned into the pGEX-1 expression vector (Pharmacia,
10 Piscataway, NJ). Inserts cloned into pGEX-1 were expressed as a fusion protein with the protein Sj26, which is a 26 Kdal Glutathione S-transferase from the parasite Schistosoma japonicum. Limited paneling of pGEX-MTA-1 against sera from HTLV-I or HTLV-II infected
15 individuals has revealed no significant difference between the reactivity of pGEX-MTA-1 vs β -gal-MTA-1.

C. Peptide Purification

The recombinant peptide is purified by standard protein purification procedures which may include differential precipitation, molecular sieve chromatography, ion-exchange chromatography, isoelectric focusing, gel electrophoresis and affinity chromatography. In the case of a fused protein, such as the β -galactosidase fused protein prepared as above, the protein isolation techniques
25 which are used can be adapted from those used in isolation of the native protein. Thus, for isolation of a β -galactosidase fusion protein, the protein can be isolated readily by simple affinity chromatography, by passing the cell lysis material over a solid support having surface-bound anti- β -galactosidase antibody. This approach is
30 used in Example 4.

II. Peptide Immunoreactivity With .5 α MAB

The invention also includes, in another aspect, a method of detecting HTLV-I positive human sera, by reacting sera with a peptide antigen which is immunoreactive with the HTLV-I Mab produced by ATCC cell line HB8755, i.e., the .5 α Mab. The presence of HTLV-I specific antibodies in sera is detected by a reporter-labeled anti-human antibody, as described in Example 7.

10

A. HTLV-I Derived Peptides

In one embodiment, the peptides contain the immunogenic region from the 42-amino acid overlap region from above-described MTA-1, MTA-4, and MTA-5 HTLV-I peptides. These peptide antigens were further characterized to confirm the location of the immunoreactive region in the 42 amino acid sequence overlap region. The location of the immunoreactive region in the C-terminal portion of the overlap region was suggested by two lines of evidence. First, the .5 α Mab was reported to react specifically with the HTLV-I envelop protein, i.e., no reaction was observed with HTLV-II or HTLV-III (HIV-1) envelop proteins. It has since been confirmed by the applicants and their co-workers that the gp46 peptide antigens MTA-1 and MTA-4 described above are reactive with HTLV-I, but not HTLV-II or HTLV-III antisera (Lipka).

Secondly, a comparison of the amino acid sequence of MTA-1 peptide with the corresponding region in the HTLV-II gp46 protein (Figure 3) shows substantially

greater homology in the N-terminal half of the peptide than in the C-terminal half (the center region of the HTLV-I and HTLV-II sequences seen in Figure 3). This would indicate that the greatest differences in antigenicity would be found in the C-terminal half of the peptide region.

This was further confirmed by antigenicity plots of the two corresponding peptide regions, shown in Figures 4A and 4B for HTLV-I and HTLV-II peptides, respectively. The antigenicity plots were generated by a standard hydrophobicity program "Antigen" in PC Gene from Intelligenetics (Palo Alto, CA). As seen, the two plots are substantially overlapping in residues 3-28, but diverge markedly in residues 28-40. The divergent residues include the HTLV-I sequence Leu-Pro-His-Ser-Asn-Leu-Asp-His-Ile-Leu-Glu-Pro-Ser.

A number of peptide antigens which include the C-terminal region just indicated were prepared and tested for binding to .5 α Mab, and to HTLV-I and HTLV-II antisera. The sequences of several of these peptides are indicated in the upper portion of Figure 3, along with the sequences of the above MTA-1, MTA-4, and MTA-5 peptide antigens. The peptides were prepared by solid-phase synthetic methods, according to standard procedures. Briefly, N-alpha-protected amino acid anhydrides were prepared in crystallized form and used for successive amino acid addition at the N-terminus. At each residue addition, the growing peptide (on a solid support) was acid treated to remove the N-alpha-protective group, washed several

18

times to remove residual acid and to promote accessibility of the peptide terminus to the reaction medium. The peptide is then reacted with an activated N-protected amino acid symmetrical anhydride, and the solid support is washed.

At each residue-addition step, the amino acid addition reaction may be repeated for a total of two or three separate addition reactions, to increase the percent of growing peptide molecules which are reacted. Typically, 1-2 reaction cycles are used for the first twelve residue additions, and 2-3 reaction cycles for remaining residues. After completing the growing peptide chains, the protected peptide resin is treated with liquid hydrofluoric acid to deblock and release the peptides from the support.

The peptides were tested for specific immunoreactivity with .5 α Mab by binding competition studies, substantially as described in Example 6. The K163 peptide, which contains the 18 C-terminal residues of MTA-4 or MTA-1, strongly inhibits binding of .5 α Mab to MTA-4. No binding interference, however, was observed with peptide K162, which contains only the 11 C-terminal residues of MTA-4. Peptide K164, which contains the 6 C-terminal residues of MTA-4 and an additional C-terminal 13 residues, weakly inhibited binding between .5 α Mab and MTA-4 or MTA-1.

These results indicate that the most potent immunoreactive region in the gp46 peptide for the .5 α Mab is in a region which includes peptide K163, consistent with the divergence in sequence homology and anti-

genicity plots between HTLV-I and HTLV-II sequences in this region. The weak binding of .5 α Mab to the K164 peptide may indicate that the epitope of interest in the His-Ile-Leu-Glu-Pro-Ser-His-Ile-Leu region of overlap between K163 and K164, where adjacent N-terminal or C-terminal sequences are required for antigen presentation, or may indicate that the K164 peptide contains an additional epitopic region which is weakly immunoreactive with the .5 α Mab.

10 The peptides were also examined for their ability to inhibit binding of antisera from HTLV-I infected patients to MTA-4. In general, it was found that the ability of any particular peptide to inhibit binding of .5 α Mab to MTA-4 paralleled its ability to either

15 bind to HTLV-I antisera in an ELISA binding protocol (Example 6B) or to inhibit binding of human HTLV-I antisera to MTA-4 or MTA-1 in a Western blot assay (Example 8C). Thus, peptide K162 did not react with any HTLV-I sera in the ELISA protocol and did not

20 inhibit binding of J-254 sera to MTA-1 or MTA-4.

B. Random-Sequence Peptides

In another embodiment, the .5 α Mab-reactive peptide for use in the method is prepared by selection

25 of random-sequence peptides. Recently, it has been demonstrated that antibodies directed against specific short (5-10 residues) peptides can be used to screen libraries of randomly generated peptides for immunoreactive species. (Scott; Cwirla et al). Such a

30 strategy is exploited herein to identify novel

20

sequences which are immunoreactive with the .5 α monoclonal antibody.

In the preferred method, approximately 10^8 novel heptapeptides are generated through construction of an epitope library using the filamentous phage fUSE5 as a vector. Other filamentous phage vectors are considered to be equally efficacious in developing such a library.

Figure 5 shows schematically the sequence of steps necessary to generate and screen a fUSE5 filamentous phage epitopic library. Briefly, fUSE5 RF DNA is subjected to digestion with restriction endonuclease SfiI to create an insertion site for insertion of foreign DNA. A synthetic (15+3m) base pair (bp) BglI DNA fragment is prepared which contains a degenerate sequence of the form (NNK)m, where N represents A, G, C, or T; K represents G or T; and m can vary from 2 to 15. In the preferred embodiment of the invention, m ranges from 5-10 and the bases are randomly added in single addition events to the template primer. An alternative method of achieving random addition of codons coding for the twenty amino acids is to randomly attach trinucleotide codons representing each amino acid to the template primer.

Following ligation of the insert to the cloning vector, amplification of the filamentous phage vector is achieved by transfection of E. coli cells. Successful transfection is measured by the presence of vector borne markers. In the preferred embodiment of the invention, this marker is tetracycline resistance.

Recombinant phage are then isolated from bacterial cells. Phage bearing sequences of interest are isolated by an antibody panning method in which phage are incubated with the .5 α Mab or its Fab fragment. 5 Biotinylated second antibody (goat anti-human IgG) is then added, and complexes containing biotinylated second antibody, the .5 α Mab and immunoreactive peptide bearing phage are separated from unreacted antibodies and phage by adhesion onto a streptavidin 10 coated plate. Phage bearing immunoreactive sequences are then eluted, and their DNA sequences are determined.

Foreign DNA sequences present in the filamentous phage fusion protein pIII determine the sequence of 15 the immunoreactive peptide. Peptides discovered to be immunoreactive through this procedure can then be synthesized by standard peptide synthetic methods and prepared as immunogens by conjugation to an appropriate peptide carrier.

20

III. HTLV-II Peptide Antigens

This section describes the identification and cloning of HTLV-II peptides which are specifically immunoreactive with HTLV-II antisera. The peptides 25 are derived from the HTLV-II gp46 envelop protein region which is homologous to the above described MTA-1 peptide from the HTLV-1 gp46 region.

An HTLV-II peptide designated GH2-K15 (Figure 3) corresponding to the HTLV-I peptide MTA-1 was prepared 30 by cloning of an HTLV-II coding sequence corresponding

to the desired peptide sequence. A 147 base pair (bp) HTLV-II DNA fragment corresponding to nucleotides 5648 to 5794 of the HTLV-II genome (Figure 6) was originally amplified from the HTLV-II clone pM04 (which contains the majority of the HTLV-II genome cloned into the BamH I site of the plasmid pBR322) by use of the polymerase chain reaction (PCR) procedure (Perkin-Elmer/Cetus GeneAmp kit).

The forward direction and reverse primers are indicated in Figure 6. The amplified DNA was ligated into the EcoR I site of λ gt11 phage vector, yielding the clone as3K15 which contains a 147 HTLV-II DNA insert into the -galactosidase gene of the λ gt11. The recombinant phage was used to transfect E. coli strain BNN103. Details are given in Example 5.

In a preliminary experiment, sera from approximately 200 individuals with PCR-confirmed HTLV-I or HTLV-II infection, as well as sera from approximately 150 uninfected individuals were paneled against the GH2-K15 antigen. 98% of the sera from HTLV-II infected individuals reacted with GH2-K15. None of either the HTLV-I infected sera or the uninfected sera reacted with GH2-K15. The screening results demonstrate that the GH2-K15 peptide is specifically immunoreactive with HTLV-II positive sera.

Several smaller peptides contained with the GH2-K15 amino acid sequence were prepared by recombinant methods, as outlined in Section I. Briefly, the peptides were prepared by PCR amplification of HTLV-II genomic DNA, using PCR primers designed to promote

amplification of the sequences indicated, as detailed in Example 5. Five of these peptides, designated (GH2-) K14, K16, K24, K35, and K34 have the sequences shown in Figure 3.

5 The recombinant HTLV-II peptides described above were immunoscreened against several HTLV-II and HTLV-I in an ELISA format, as described in Example 8. The results are shown in Table 1. All λ gt11 HTLV-II clones except for GH2-K16 were recognized by at least 1 out
10 of the 6 HTLV-II sera tested. GH2-K16, the sole non-reactive clone, is missing the carboxyl terminal 22 amino acids that are included in GH2-K15. All the other clones tested contain at least the 17 amino acids Ser-Pro-Pro-Leu-Val-His-Asp-Ser-Asp-Leu-Glu-His-
15 Val-Leu-Thr-Pro-Ser that are present in peptide K125.

Also as seen in Table 1, none of the tested peptides reacted with any of the HTLV-I sera, nor with the .5 α Mab.

Three of the original HTLV-II clones, GH2-K15,
20 GH2-K35, and GH2-K16 have been cloned into the pGEX-1 expression vector. Recombinant protein expressed by the 3 pGEX-1 HTLV-II clones GH2-K15, GH2-K25, and GH2-K35 have all been recognized by the J-317 HTLV-II serum.

24

Table 1

				-----HTLV-II ANTIGENS-----					
	SERUM	VIRUS	N	K15	K14	K16	K24	K34	K35
5	J-115	II	2	+/-	-	-	-	-	-
	J-127	II	2	-	-	-	-	-	-
	J-289	II	2	-	-	-	-	-	-
	J-309	II	2	-	-	-	-	-	-
	J-263	II	3	+/-	-	-	+	-	-
10	J-317	II	2	++	+	-	++	+	+
	J-103	I	2	-	-	-	-	-	-
	J-108	I	2	-	-	-	-	-	-
	J-183	I	2	-	-	-	-	-	-
15	.5α Mab	I	1	-	-	-	-	-	-

A number of peptide antigens which contain amino acid sequences within the K15 sequence were prepared by solid-phase methods, as outlined in Section III above.

20 The sequences of five of these peptides, designated (GH2-) K169, K170, K125, K126, and K128 are shown in Figure 3. The peptides were tested for immunoreactivity with several HTLV-I and HTLV-II positive sera, by an ELISA method, and some of the peptides were also examined
25 for their ability to inhibit HTLV-II antibody binding to the K15 antigen.

The K125 peptide was recognized by multiple HTLV-II sera when assayed by ELISA. In one experiment 6 out of 12 HTLV-II sera were able to bind efficiently to K125.
30 In the same experiment 0 out of 7 HTLV-I sera bound peptide K125. The K125 peptide also inhibited the binding of a strongly reactive HTLV-II sera, J-317, to

Western blotted GH2-K15. The ability of sera J-317 to bind GH2-K15 is not affected by incubation with the HTLV-I peptide K163 or the HTLV-II peptide K128.

5 The HTLV-II peptide K170 is recognized by multiple HTLV-II sera in an ELISA based assay, and not recognized by HTLV-I sera in the same assay. The K169 peptide is not recognized by HTLV-II sera in an ELISA based assay.

10 Data from both the analysis of HTLV-II recombinant antigens and the synthetic HTLV-II peptides indicate that the HTLV-II specific epitope is contained in the 17 amino acid sequence Ser-Pro-Pro-Leu-Val-His-Asp-Ser-Asp-Leu-Glu-His-Val-Leu-Thr-Pro-Ser in the GH2-K15 peptide. Data obtained by extensive paneling of the HTLV-I antigens MTA-1 and MTA-4, discussed above, would suggest that the
15 6 final amino acids of GH2-K15, Thr-Ser-Trp-Thr-Thr-Lys, may also contribute to the epitope recognized by HTLV-II antisera.

IV. HTLV-I and HTLV-II Diagnostic Methods

20 Three basic types of diagnostic applications of the HTLV-I and HTLV-II peptide antigens of the invention will be described. The first is based on inhibition of complement-mediated, antibody-dependent cytotoxicity by the peptide. In this method, serum from a test individual is
25 reacted with HTLV-I or HTLV-II infected T-cell clones in the presence of complement. The presence of anti-HTLV-I or anti-HTLV-II antibody is evidenced by cell lysis, as judged, for example, by trypan blue dye exclusion.

30 Where cell lysis is observed, the specificity of the anti-HTLV-I antibody for the HTLV-I peptide is demonstrated by first reacting the serum with excess HTLV-I or HTLV-II peptide, then mixing the serum with cells in the

presence of complement. The presence of HTLV-I or HTLV-II antibody is indicated by a substantial decrease in cell lysis. This method is described in Example 6A.

5 The method can also be used to quantitate the antibody titer in the analyte serum, by titrating the serum with increasing amounts of peptide, and determining the peptide concentration where a noticeable effect on the extent of cell lysis is first observed.

10 The second general assay type is an enzyme-immunoassay for screening human sera for HTLV-I or HTLV-II infection. In this assay format, a solid phase reagent having surface-bound HTLV-I or HTLV-II gp46 peptide antigen is reacted with analyte serum, under conditions which allow antibody binding to the peptide on the reagent.

15 After washing the reagent to remove unbound serum components, the reagent is reacted with an enzyme-labeled anti-human antibody, to bind enzyme to the reagent in proportion to the amount of bound anti-HTLV-I antibody on the solid support. The reagent is again washed, to remove unbound antibody, and the amount of enzyme associated with the reagent is determined. One exemplary method, employing an anti-human antibody labeled with alkaline phosphatase, is detailed in Example 7 for a direct HTLV-I screening assay. The enzyme-labeled antibody, and reagents required for enzyme detection, are

20 also referred to herein as reporter means for detecting the presence of human antibody bound to the peptide antigen on the solid support.

30 The solid surface reagent in the above assay is prepared by known techniques for attaching protein material to solid support material, such as polymeric beads, dip sticks, or filter material. These attachment methods

generally include non-specific adsorption of the protein to the support (as in the filter support described in Example 8) or the covalent attachment of the protein, typically through a free amine group, to a chemically reactive group on the solid support, such as an activated carboxyl, hydroxyl, or aldehyde group.

The third general assay type is Western blot assay for use in confirming HTLV-I or HTLV-II antisera. This assay format includes, in addition to the gp46 peptide antigen of the invention, one or more confirmatory HTLV-I or HTLV-II antigens that are effective to detect HTLV-I or HTLV-II antisera. In one preferred format, the confirmatory peptides include the p24 gag protein from HTLV-I viral lysate, and a p21E recombinant envelop protein containing a large portion of the HTLV-I gp21 envelop protein (Samuel, 1984, 1985). The p24 lysate proteins picks up most, but not all HTLV-I and HTLV-II positive sera. The p21E recombinant peptide picks up virtually all HTLV-I and HTLV-II, but also gives some false positives. This modified Western blot assay has been reported by the applicants and co-workers (Lipka). Details of the blot assay procedure are given in Example 8.

As has been described, and as is detailed in Example 8, the modified Western blot format picked up all HTLV-I and HTLV-II positive sera tested (a panel of 95), as evidence by immunoreaction with viral lysate protein p24 and recombinant protein p21E. In addition, the MTA-4 peptide was immunoreactive with confirmed HTLV-I sera only. The modified blot assay thus can be used to confirm HTLV-I or HTLV-II antisera, and to distinguish

the two types of HTLV virus by selective immunoreaction with the peptide of the invention.

In another embodiment of the Western blot assay, the HTLV-I peptide antigen is replaced by the HTLV-II gp peptide antigen described in Section III. In this format, the HTLV-I viral lysate proteins and p21E recombinant protein provide confirmation of HTLV-I or HTLV-II antisera, as above. The HTLV-II specific peptide will pick up HTLV-II, but not HTLV-I antisera, and thus provides a positive confirmation of HTLV-II antisera.

The two formats can be combined to include both HTLV-I and HTLV-II specific peptide antigens, to give positive confirmation of either HTLV antisera.

15 V. Vaccine Compositions

Also included in the invention is a vaccine composition containing an HTLV-I gp46 peptide and a antigen carrier, such as an immunogenic protein, to which the antigen peptide is bound. The peptide contains an immunogenic region formed by the above 42- or 47-amino acid overlap of MTA-1, MTA-4, and MTA-5 peptides described in Section I, which is immunoreactive with anti-HTLV-I .5 α Mab, i.e., the antibody derived from ATCC cell line HB8755. More specifically, the peptide contains the immunogenic region of the peptide sequence Thr-Ala-Pro-Pro-Leu-Leu-Pro-His-Ser-Asn-Leu-Asp-His-Ile-Leu-Glu-Pro-Ser. Since the .5 α Mab is a neutralizing antibody, the antibody raised by the peptide is expected to be a neutralizing antibody.

30 The vaccine composition may alternatively include the HTLV-II gp46 peptide containing the HTLV-II

specific immunogenic region formed by the amino acid sequence Met-Thr-Leu-Leu-Val-Asp-Ala-Pro-Gly-Tyr-Asp-Pro-Leu-Trp-Phe-Ile-Thr-Ser-Glu-Pro-Thr-Gln-Pro-Pro-Pro-Thr-Ser-Pro-Pro-Leu-Val-His-Asp-Ser-Asp-Leu-Glu-
5 His-Val-Leu-Thr-Pro-Ser-Thr-Ser-Trp-Thr-Thr-Lys, and preferably formed by the amino acid sequence Ser-Pro-Pro-Leu-Val-His-Asp-Ser-Asp-Leu-Glu-His-Val-Leu-Thr-Pro-Ser-Thr-Ser-Trp-Thr-Thr-Lys, or Ser-Pro-Pro-Leu-Val-His-Asp-Ser-Asp-Leu-Glu-His-Val-Leu-Thr-Pro-Ser.

10 Particularly useful protein carriers for the peptide(s) include keyhole limpet hemocyanin (KLH), tetanus toxoid, poly-l-(Lys:Glu), peanut agglutinin, poly-D-lysine, diphtheria toxoid, ovalbumin, soybean agglutinin, bovine serum albumin (BSA), human serum
15 albumin, and the like.

The immunogenic peptide(s) may be conjugated to the carrier by a variety of known methods, including chemical derivatization and by genetic engineering techniques. Such latter technique is disclosed in
20 more detail by Gerald Quinnan, "Proceedings of a Workshop," November 13-14, 1984. Vaccines and inocula of the present invention may be administered by injection, usually intramuscularly or subcutaneously, orally by means of an enteric capsule or tablet, as a suppository, as a nasal spray, and by other suitable
25 routes of administration. For a human patient, a suitable dose of the polypeptide depends, in part, upon the chosen route of administration and a number of other factors. Included among those factors are
30 the body weight of the mammal to be immunized, the

30

carrier when used, the adjuvant when used, and the number of inoculations desired to be used.

Individual inoculations for a human patient typically contain unit doses of about 10 micrograms to about 100 milligrams of polypeptide, exclusive of any carrier to which the polypeptide may be linked. If desired, a series of doses may be administered over a period of time for optimum immunity. Unit dosage forms of the vaccine can also be provided, if desired, containing the aforementioned amounts of the polypeptide.

In any event, the immunogen contained in a vaccine or an inoculum is present in an "effective amount," which amount depends upon a variety of factors as is well known in the immunological arts, e.g., the body weight of the mammal to be immunized, the carrier moiety used, the adjuvant used, the duration of protection sought, and the desired immunization protocol.

The following examples illustrate various aspects of the invention, but are in no way intended to limit the scope thereof.

Materials

The materials used in the following Examples were as follows:

Enzymes: DNAase I and alkaline phosphatase were obtained by Boehringer Mannheim Biochemicals (BMB, Indianapolis, IN); EcoRI, EcoRI methylase, DNA ligase, and Polymerase I, from New England Biolabs (NEB, Beverly, MA); and RNase was obtained from Sigma (St. Louis, MO).

Other reagents: EcoRI linkers were obtained from NEB; and nitro blue tetrazolium (NBT), 5-bromo-4-chloro-3-indolyl phosphate (BCIP), 5-bromo-4-chloro-3-indolyl- -D-galactopyranoside (X-gal) and isopropyl -
5 D-thiogalactopyranoside (IPTG) were obtained from Sigma.

Example 1

Preparation of an HTLV-I Genomic Library Source of Genomic Material

10 Bacteriophage containing a full-copy DNA insert derived from the HTLV-I genome was obtained from Drs. R.C. Gallo and F. Wong-Staal of the Laboratory of Tumor Cell Biology, National Institutes of Health (Bethesda, MD). The bacteriophage was digested to
15 completion with SacI, releasing the viral genome insert. The digested material was electrophoresed on standard 1% agarose gel, and the 9.5 kilobase fragment obtained by electroelution was extracted with phenol/chloroform before ethanol precipitation.

20 The purified genomic DNA was suspended in a standard digest buffer (0.5M Tris HCl, pH 7.5; 1 mg/ml BSA; 10mM MnCl₂) to a concentration of about 1 mg/ml, and digested with DNAase I at room temperature for about 5 minutes. These reaction conditions were
25 determined from a prior calibration study, in which the incubation time required to produce predominantly 100-300 basepair fragments was determined. The material was extracted with phenol/chloroform before ethanol precipitation.

30 The genomic fragments from above were blunt-ended with DNA Pol I under standard conditions (Huynh), then

extracted with phenol/chloroform and precipitated with ethanol. The blunt-ended material was ligated with EcoRI linkers, under standard conditions (Maniatis, pp. 396-397), then digested with EcoRI to remove
5 redundant linker ends. The material was then agarose-gel-fractionated to remove non-ligated linkers and to size-select (see below).

The resultant fragments from the previous step were analyzed by electrophoresis (5-10V/cm) on 1.2%
10 agarose gel, using X174/HaeIII and /HindIII size markers. The 100-300 bp fraction was eluted onto NA45 strips (Schleicher and Schuell), which were then placed into 1.5 ml microtubes with eluting solution (1 M NaCl, 50 mM arginine, pH 9.0), and incubated at 67°C
15 for 30-60 minutes. The DNA, now in solution, was extracted with phenol/chloroform and precipitated with ethanol. The pellet was resuspended in 20 µl TE (0.01 M Tris HCl, pH 7.5, 0.001 M EDTA).

gt11 phage vector (Huynh) was obtained from Promega
20 Biotec (Madison, WI). This cloning vector has a unique EcoRI site 53 base pairs upstream from the β-galactosidase translation termination codon. The genomic fragments from above were introduced into the EcoRI site by mixing 0.5 -1.0 µg EcoRI-cleaved gt11, 0.5-3 µl of the
25 above HTLV--I genomic fragments, 0.5 µl 10X ligation buffer (above), 0.5 µl ligase (200 units), and distilled water to 5 µl. The mixture was incubated overnight at 14°C, followed by *in vitro* packaging, according to standard methods (Maniatis, pp. 256-268).

30 The packaged phage were used to infect E. coli, strain KM392, obtained from Dr. Kevin Moore, DNAX (Palo

33

Alto, CA). Alternatively, E. coli strain Y1090, available from the American Type Culture Collection (ATCC #37197), could be used. The infected bacteria were plated and the resultant colonies were checked for loss of β -galactosidase activity (clear plaques) in the presence of X-gal using a standard X-gal substrate plaque assay method (Maniatis). Table 2 below shows the number of recombinant (clear) plaques obtained with insertion of the EcoRI-ended HTLV-I fragments (row 1). An EcoRI linker control (row 2) and vector alone (row 3) were also run. As seen, about 50% of the phage plaques showed loss of enzyme (recombination). The background levels either in the presence or absence of EcoRI linkers were less than 15%, indicating the successful generation of an HTLV-I epitope library. The phage libraries contained about 10^6 plaque-forming units (pfu)/ml.

Table 2

	<u>Insert</u>	<u>Vector</u>	<u>Clear/Total</u>	<u>%Rec</u>
20	1. <u>SacI</u> in 3.25 μ l	1 μ l	100/200	50
25	2. <u>EcoRI</u> linker 3.25 μ l	1 μ l	25/178	14
	3. Control	1 μ l	50/400	13

30

Example 2Screening for gp46 Coding Inserts

Purified .5 antibody derived from a human cell line (ATCC #C8755) was provided by Dr. Samuel Broder of the
5 National Cancer Institute, National Institutes of Health (Bethesda, MD). Mouse anti-human IgG antibody covalently derivatized with alkaline phosphatase was obtained from Promega Biotec (Madison, WI).

A lawn of KM392 cells infected with about 10^4 pfu of
10 the phage stock from Example 1 was prepared on a 150 mm plate, and incubated, inverted, for 5-8 hours at 37°C. The lawn was overlaid with a nitrocellulose sheet, causing transfer of HTLV-I recombinant protein from the plaques to the paper. The plate and filter were indexed
15 for matching corresponding plate and filter positions.

The filter was washed twice in TBST buffer (10 mM Tris, pH 8.0, 150 mM NaCl, 0.5% Tween 20), blocked with AIB (TBST buffer with 1% gelatin), washed again in TBST, and incubated overnight after addition of .5 monoclonal
20 antibody (diluted to 1-2 µg/ml in AIB, 12-15 ml/plate). The sheet was washed twice in TBST, then contacted with enzyme-labeled anti-human antibody, to attach the labeled antibody at filter sites containing antigen recognized by the .5 antibody. After a final washing, the filter was
25 developed in a substrate medium containing 33 µl NBT (50 mg/ml stock solution maintained at 5°C) in 5 ml of alkaline phosphatase buffer (100 mM Tris, 9.5, 100 mM NaCl, 5 mM MgCl₂). Reacted substrate appeared at points of antigen production, as recognized by the 0.5α Mab.

30 The areas of antigen production determined in the previous step were replated at about 100-200 pfu on an 82 mm plate. The above steps, beginning with a 5-8 hour

35

incubation, through NBT/BCIP development, were repeated in order to identify plaques which secreted an antigen capable of reacting with the .5 Mab. The identified plaques were picked and eluted in phage buffer (Maniatis, p. 443). Three of the recombinant phage plaques which secreted an antibody-reactive peptide were selected for sequencing analysis, according to the procedures in Example 3. The corresponding infected phage has been designated MTA-4, MTA-1, and MTA5.

10

Example 3

Phage Purification and DNA Extraction

Phages MTA-4, MTA-1, and MTA-5 were isolated from the plate cultures of the infected E. coli Y1088 bacteria. These cells are available from the ATCC (ATCC #31195). The phage was collected by addition of phage-dilution buffer (Maniatis) late material was purified from bacterial debris by low-speed centrifugation, and the supernatant was poured into SW 27 tubes. RNase and DNase were each added to a concentration of 1µg/ml each from stock solutions of 1 mg/ml. The sample was incubated for 30 minutes at 37°C, and an equal volume of a polyethylene glycol (PEG), 5.8 g NaCl, 2.0g MgSO₄•7H₂O, 1M Tris Cl, pH 7.5, and 2% gelatin was added. The sample was placed in an ice bath for 1 hour to allow the phage particles to form a precipitate, which was then isolated by centrifugation at 10k for about 20 minutes at 4°C.

The supernatant was decanted, and the pellet was re-suspended in 0.6 ml PDB buffer (5.8 g NaCl, 2.0 g MgSO₄•7H₂O, 50 ml 1M Tris Cl, pH 7.5, and 5 ml 2% gelatin) and transferred to .5 ml polypropylene microtubes. 5 µl

36

10% SDS, 5 μ l 0.5M EDTA, and 2.5 μ l proteinase K (20 mg/ml) were added, and the samples were incubated at 50°C for 15 minutes.

5 The detergent and enzyme-treated material was extracted with an equal volume of phenol/chloroform, and centrifuged to ensure separation of the phases. The aqueous phase was transferred to a new tube, and the extraction/centrifugation procedure was repeated with a mixture of chloroform and isoamyl alcohol. An equal
10 volume of isopropanol was added, and the same was inverted several times to mix, and cooled to -70°C for 20 minutes. The sample was centrifuged for 5 minutes and the supernatant was decanted. The pellet was washed in 70% ethanol, briefly dried in a 37°C heat block, and re-
15 suspended in 100 μ l TE buffer, pH 7.5.

The isolated phage DNA was digested with KpnI and SacI and then combined with KpnI/SacI cut plasmid vector pGEM-3 (Promega Biotec) to isolate a plasmid recombinant with the insert of interest. The HTLV-I insert was then
20 sequenced using the standard dideoxy sequencing procedure and forward and reverse primers for λ gt10 sequences flanking the EcoRI insertion site.

The figure shows the coding sequence and corresponding amino acid sequence of a portion of the fused protein
25 formed by the above methods, for each of the three fused peptides examined. A terminal G base of the β -gal gene and the adjacent CC bases of the env gene contributed by each of the three insert sequences yield a GCC (Ala) codon, replacing the Ser codon which normally occurs at
30 that codon position of all three env inserts. As shown, the insert in the MTA-4 includes a 225 base pair sequence

extending from base 5564 to 5790 of the HTLV-I coding region. The insert of the MTA5 phage begins at base 5664, and extends to base 5895. The 231 basepair sequence covers amino acids 162 to 240 of the gp46 protein.

5 The region of insert overlap, from 5664 to 5790, includes the 42 amino acid sequence from amino acids 162 to 203 of the native gp46 protein.

Example 4

10 Isolation of HTLV-I Peptide Antigens

A. Construction of Lysogens

EcoRI, strain C600, was obtained from Dr. R. Davis, Stanford University (Stanford, CA). Alternatively, EcoRI Y1089 (ATCC #37196) can be used. A 1 ml saturated, over-
15 night culture of the cells was infected with one of the three phages from Example 3 by adsorbing 10 µl of eluted plaque stock to 50 µl of overnight bacterial culture. The infected bacteria were spread only LB agar plates (Maniatis, p. 440) and incubated at 32°C. The individual
20 colonies were picked with sterile toothpicks onto corresponding grids on two separate plates. One of the plates was incubated at 32°C, and the other at 42°C. Cells that grew at the lower temperature (indicating a lysogenic state produced by the presence of the phage repressor
25 protein) but not at the higher temperature (because of cell lysis) were assumed to be lysogenic. Many lysogenic colonies from each of the three phage types were found.

B. Recombinant Antigen Induction from Lysogens

This Example describes induction of a recombinant protein containing the HTLV-I epitope from the λ gt11 lysogens prepared in Example 4 with the MTA-4 phage. As indicated above, the antigen is produced in the form of a -galactosidase fusion protein which also contains an N-terminal portion of the phage -gal protein.

A superbroth was prepared containing 35 g bacto-tryptone, 2 g bacto-yeast extract, 5 g NaCl, and 5 ml 1N NaOH in 1 l dH₂O. 500 ml of the superbroth were inoculated 1:100 with a saturated overnight culture of the EcoRI λ gt11 lysogens prepared in the previous example. The culture was incubated to $A_{600} \approx 0.4-0.5$ with vigorous aeration.

In order to maximize protein production, the temperature of the culture was raised to 43-44°C, thereby inactivating the temperature-sensitive -galactosidase repressor gene. The temperature was maintained at 43°C with a 65°C water bath for 15 minutes with aeration. IPTG, which induces -galactosidase expression by competitively binding to the -galactosidase repressor, was added to the broth to 2 mM to further increase protein production. The culture was returned to the 38°C shaker for about an hour. The cells were then pelleted at 6,000 x g for 15 minutes at 37°C, resuspended in lysis buffer (10 mM Tris, pH 7.4, 2% Triton X-100, 1% aprotinin, and 50 μ g PMSF) and immediately plunged into liquid N₂. Lysis was completed upon thawing of the frozen samples.

C. Purification of Fusion Protein

The cell lysate obtained in the previous Example was thawed and warmed to 37°C. 10 µl DNase (1 µg/ml) was added, and the mixture incubated until the viscosity decreased. The lysate was quickly chilled on ice, clarified at 4°C for 5 minutes in a microfuge, and loaded onto a 6 ml column of anti-β-galactosidase coupled to Sepharose 4B (Pharmacia). The column was allowed to equilibrate 1-2 hour, and washed with 7 volumes (column volumes) of TX buffer (10 mM Tris, pH 8.0, 2% Triton X-100, 50 µg/ml PMSF), followed by 2 volumes of 5mM 3,5-diiodosalicylic acid in TX buffer. Fusion protein was then eluted from the column with 35 mM 3,5-diiodosalicylic acid in TX buffer. The majority of protein was eluted in the first 3-4 volumes, and removal was substantially complete after 7 volumes.

The eluted samples were desalted and concentrated using Amicon filters (Danvers, MA).

20

Example 5

Preparing HTLV-II Antigens

A. Synthesis and Cloning of HTLV-II DNA Sequences

The polymerase chain reaction (PCR) procedure was used to generate HTLV-II DNA sequences for cloning. Six 30 bp DNA primers were synthesized. All 6 primers had 3 bp of gtl1 sequence followed by an EcoR I site at their 5' ends. This was followed by 21 bp of HTLV-II sequences. The 3 forward direction primers contained HTLV-II sequences corresponding to nucleotides 5648 - 5668, 5687 - 5707, and 5726 - 5745. The 3 reverse direction primers contained HTLV-II sequences

40

corresponding to nucleotides 5794 - 5774, 5776 - 5756, and 5728 - 5708.

PCR was performed according to the manufacturers instructions (Perkin Elmer/Cetus), and all PCR reactions
5 contained 2 ng of the above HTLV clone as template and 1 μ M of the appropriate PCR primers. PCR amplification was carried out for 25 cycles. Each cycle involved template denaturation for 1 minute at 94 deg.C, annealing of primer to template for 2 minutes at 50 deg.C, followed by
10 primer extension for 2 minutes at 72 deg.C. Afterwards the amplified DNA was purified and then digested to completion with EcoRI. The digested DNAs were then ligated into the EcoRI site of lambda gt11. The recombinant phage DNAs were then packaged and the
15 frequency of non-recombinant phage was determined by plating in the presence of 5-bromo-4-chloro-3-indolyl β -D-galactopyranoside.

The ratio of recombinant to non-recombinant phage was about 50/1. Multiple isolated plaques from each of
20 the 6 recombinant phage clones were picked and subsequently screened using PCR with lambda gt11 flanking primers 11F and 11R, and/or the HTLV-II plaques described above. Clones containing correctly sized and orientated inserts were then amplified and used in subsequent
25 immunoscreening assays. The EcoR I fragment from 3 of the clones GH2-K15, GH2-K16 and GH2-K35 were subsequently subcloned into the pGEX-1 plasmid and DNA sequenced. The sequences obtained perfectly matched the reported sequence for the desired region of HTLV-II (Shimotokno).
30 B. Immunological Analysis of HTLV-II Clones

Recombinant phage was mixed 1/1 with wild type gt11 and used to infect E. coli strain KM-392. After allowing

the phage to grow for ~5 hours expressed proteins were bound to nitrocellulose filters overnight. Filters were subsequently washed 3X with TBS (0.5 M NaCl, 20 mM Tris Ph 8.0), cut into sections, and blocked using TBS plus 1% Gelatin. Filter sections were then incubated overnight with 1st stage antibody, usually sera from HTLV-I or HTLV-II infected individuals diluted 1/100 in TBS plus gelatin. After washing with TBS, the filters were incubated with alkaline phosphatase conjugated goat anti human sera for at least 1 hour. The filters were washed with TBS and bound antibody was then detected by incubating the filters in a solution of nitroblue tetrazolium chloride and 5-bromo-4-chloro 3-indolylphosphate. A particular sera was scored as positive if plaques derived from the recombinant phage could clearly be distinguished from plaques of wild type gt11.

C. Expression and Purification of Recombinant Antigen
β-galactosidase fusion proteins were expressed by first generating lysogens. Recombinant gt11 phage was used to infect E. coli strain BNN103, and lysogens were identified by growing duplicate plates at 32°C and 42°C. The production of fusion protein was induced by raising the temperature of a log phase culture of lysogen to 42°C for 15 minutes. Isopropyl thiogalactoside was then added to a final concentration of 1.6 Mm and the cultures were grown for an additional 1 hour at 37°C. Cells were then pelleted by centrifugation at 5000 x g for 15 minutes and resuspended in 1/50th original culture volume of lysis buffer (2% Triton X-100, 1% Aprotinin, 10 mM Tris, pH 7.4). The solution was then frozen by immersion in a dry

42

ice / ethanol bath and then thawed. DNase I was added to a final concentration of 1 $\mu\text{g}/\text{ml}$ and the lysate was then incubated for 5 minutes at room temperature. Insoluble debris was then pelleted by centrifugation at $10,000 \times g$ for 10 minutes. The supernatant was then centrifuged as before. sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE, Lammeli) analysis of aliquots of the pellet and supernatant fractions indicated that GH2-K15 was found primarily in the supernatant fraction.

10 The supernatant fraction was then combined with 2 mls of Protosorb LacZ adsorbent (Promega) and incubated for 2 hours at 25°C with agitation. The column resin was then poured into a disposable column and washed with 2X with 10 mls of TX buffer (1% Aprotinin, 10 mM Tris pH 15 7.4). Bound protein was eluted 14 mls of pH 10.8 carbonate buffer and 2 ml fractions were collected into tubes already containing 1 ml of 2 M tris buffer (pH 7.5). Fractions were then concentrated using Centricon 30s (Amicon) following manufacturers instructions. The 20 fractions were washed with 2 mls of MTBS buffer (150 mM NaCl, 4 mM NaH_2PO_4 , 16 mM Na_2HPO_4 , pH 7.3) and then concentrated again.

The location, yield and purity of the purified fusion protein was determined using SDS-PAGE. After one 25 pass through the immunoaffinity column GH2-K15 recombinant antigen was ~70 % pure. Fractions containing fusion protein were pooled and aliquots of the pool were used in subsequent western blot experiments. Western blot analysis was performed essentially as described previously (Lipka). Titration experiments determined 30 that the optimum loading of purified GH2-K15 antigen was 3 μg protein/cm nitrocellulose. Peptide competition

experiments are described in Example 6C. Nitrocellulose strips containing blotted antigen were then added to the serum samples and incubated and developed as normal.

D. HTLV Antisera

5 Sera samples were obtained from HTLV-I, HTLV-II, or ELISA reactive - HTLV negative individuals. These samples were from multiple different sources and geographic areas. Many of the seropositive sera samples were also typed for HTLV-I or HTLV-II infection using PCR.
10 with strain specific DNA primers. HTLV-I sera samples included 58 PCR proven samples consisting of 45 samples from Jamaican food handlers, 2 intravenous drug users (IVDU) from the New Orleans area, and 11 northern California blood donors. In addition a total of 238
15 HTLV-I sera samples were obtained from Japan. For the Japanese samples PCR data was unavailable and the infection was typed by western blot analysis of the sera samples against HTLV-I antigens using previously described criteria (Lipka et al. JID). HTLV-II sera
20 samples included 57 PCR proven samples consisting of 6 IVDU from the New Orleans area, 24 IVDU from the northern California area, and 27 blood donors from the northern California area. HTLV negative sera included 1 Jamaican food handler, 15 California blood donors, and 29 samples
25 from Japan. PCR analysis of serum samples was performed as described (Lipka).

Example 6

Detecting Peptide Antigen Immunoreactivity

30 A. Inhibition of Cell Lysis

HUT 102-B2 cells were obtained from Dr. R.C. Gallo, LTCB, NIH. This is a long-term cultured T-cell line known to produce HTLV-I.

5 .5 α antibody (~5 μ g/ml IgG) or a control isotype matched human IgG was preincubated with MTA-4 recombinant peptide or irrelevant recombinant for 30 minutes at room temperature. 50 μ l of these mixtures was then added to 5x10⁵ HUT 02B2 cells in 96-well micro titer plates, and incubated for 30 minutes at room temperature. 30 μ l of
10 rabbit complement per well was added, and incubated 1 hour at 37°C. Cell viability was determined by microscopic examination. Cell lysis was visibly inhibited by addition of the MTA-4 peptide antigen, but not by preincubation with irrelevant recombinant peptide
15 antigen. Isotyped matched human IgG, after preincubation with either recombinant antigen or irrelevant recombinant peptide antigen, had no effect on HUT 102-B2 viability.

B. ELISA Assay

20 HTLV-I and HTLV-II peptides were examined in an ELISA assay to determine the ability of sera from HTLV infected individuals to bind to the synthetic peptides described above. Briefly, the ELISA assay involved binding a fixed amount of synthetic peptide to a microtiter plate, followed by the addition of sera from a
25 HTLV infected individual. Unbound sera was then washed away and antibody bound to the peptide was detected by a 2nd antibody. The 2nd antibody is conjugated to an enzyme that converts a colorless substrate to a colored product. The amount of colored product produced
30 indicates the amount of serum antibody which bound the peptide. The signal obtained from a particular sera against bound peptide was subtracted from the signal

45

obtained by the sera from a well which did not contain any peptide. The values obtained after subtraction of the minus peptide background had to be 2.5 times the background value to be considered positive.

5

C. Antibody Binding Inhibition

The inhibition assay involves the incubation of a large excess of a synthetic peptide with sera from an HTLV infected individual prior to placing a strip of
10 nitrocellulose which contains a HTLV-I or HTLV-II recombinant antigen blotted on to it. If the sera can bind to the peptide, the vast excess of peptide in solution with the antibody will prevent significant binding of the antibody to the relatively small amount of antigen present
15 on the nitrocellulose strip. The amount of antibody bound to the recombinant antigen on the nitrocellulose strip is determined using an enzyme conjugated second antibody in a manner analogous to that described above for the ELISA assay. Control experiments involved incubating
20 HTLV-I sera with the HTLV-II peptide K125 or HTLV-II sera with an HTLV-I peptide, and then determining the ability of the sera to recognize the appropriate recombinant antigen.

25

Example 7

EIA Assay

Purified MTA-4 peptide antigen was prepared as in Example 4, and dot blotted on nitrocellulose filters, which were then used in a solid-phase assay for
30 determination of serum antibodies in patients with T-cell leukemia (6 patients with HTLV-I infection). In each case, 0.1 ml of various serum dilutions, ranging from

46

1:100 to 1:50,000, from the test individual was added to the filter, and allowed to rest at room temperature for 30 minutes. The filter was then washed two times with TBST buffer (Example 2), and incubated with anti-human antibody conjugated with alkaline phosphatase, as in Example 2. The presence of antibody was determined by color development in NBT and BCIP, also as in Example 2.

Example 8

10 Modified Western Blot for Confirming HTLV-I Positive Sera

Recombinant MTA-1 was prepared as in Example 4. Recombinant p21E was prepared as described previously (Samuel, 1984, 1985). HTLV-II viral lysate was prepared from chronically infected cell line MT-2 (Hillcrest Biologicals, Cypress, CA). These HTLV-I antigens were combined and then separated under reducing conditions on a 11.5% acrylamide SDS/PAGE gel (Laemmli). The resolved proteins were electroblotted onto a nitrocellulose (onto a nitrocellulose membrane blocked with blotto (5% nonfat dry milk, 2.5% normal goat serum in 100 mM Tris-HCl, pH 7.4), air dried, and cut into 3 mm wide strips.

In the assay, the test strips from above were first rehydrated in TBS buffer, and the strips were incubated overnight with human test sera, diluted 1:50 in blotto. The strips were washed several times with wash buffer, then incubated for one hour with goat anti-human IgG conjugated to alkaline phosphatase (Bio-Rad, Richmond, CA). After washing, color development was achieved by incubating the strips in a substrate solution containing 5-bromo-4-chloro-3-indolyl phosphate and nitroblue tetrazolium in 100 mM Tris-HCl buffer, pH 9.5, 50 mM MgCl₂. Color development was continued until a uniform

background developed on the strip and was halted by rinsing the strips two times with de-ionized water.

A panel of HTLV-I or HTLV-II positive sera were tested. These had been previously confirmed as HTLV-I or HTLV-II positive by PCR analysis (Lipka). The results are shown in Figure 6, where panels A-G are HTLV-I antisera, and panels H-S are HTLV-2 antisera. Viral lysate protein gp24 was immunoreactive with every serum sample, as was the recombinant peptide gp21E. MTA-4 was immunoreactive with HTLV-I serum samples only.

While the invention has been described with reference to particular embodiments, methods of construction, and uses, it will be clear to those in the art that various other uses, formulations, and methods of practice are within the contemplation of the present invention.

IT IS CLAIMED:

1. A peptide antigen comprising less than about 77 amino acids derived from HTLV-I envelope protein gp46, said antigen including the immunogenic region formed by the amino acid sequence Ser-Leu-Leu-Val-Asp-Ala-Pro-Gly-Tyr-Asp-Pro-Ile-Trp-Phe-Leu-Asn-Thr-Glu-Pro-Ser-Gln-Leu-Pro-Pro-Thr-Ala-Pro-Pro-Leu-Leu-Pro-His-Ser-Asn-Leu-Asp-His-Ile-Leu-Glu-Pro-Ser which is immunoreactive with anti-HTLV-I monoclonal antibody derived from ATCC cell line HB8755.

2. The antigen of claim 1, wherein the immunogenic region is formed by the amino acid sequence which additionally includes the amino acid sequence Ile-Pro-Trp-Lys-Ser-Lys extending from the Ser terminus of the claim 1 sequence.

3. The antigen of claim 1, wherein the immunogenic region is formed by the amino acid sequence Thr-Ala-Pro-Pro-Leu-Leu-Pro-His-Ser-Asn-Leu-Asp-His-Ile-Leu-Glu-Pro-Ser.

4. A kit for detecting the presence of HTLV-I infection in a human serum, comprising a solid support with an immobilized gp46 peptide antigen comprising less than about 77 amino acids derived from HTLV-I envelope protein gp46, said antigen including the immunogenic region formed by the amino acid sequence Ser-Leu-Leu-Val-Asp-Ala-Pro-Gly-Tyr-Asp-Pro-Ile-Trp-Phe-Leu-Asn-Thr-Glu-Pro-Ser-Gln-Leu-Pro-Pro-Thr-Ala-Pro-Pro-Leu-Leu-Pro-His-Ser-Asn-Leu-Asp-His-Ile-Leu-Glu-Pro-Ser

which is immunoreactive with anti-HTLV-I monoclonal antibody derived from ATCC cell line HB8755, and constructed to permit serum antibodies to contact with the peptide antigen, when such serum is applied to the solid support,
5 and

reporter means for detecting the presence of human antibodies bound to said support.

5. The kit of claim 4, wherein the gp46 antigen's
10 immunogenic region additionally includes the amino acid sequence Ile-Pro-Trp-Lys-Ser-Lys extending from the Ser terminus of the claim 1 sequence.

6. The kit of claim 4, wherein the immunogenic
15 region is formed by the amino acid sequence Thr-Ala-Pro-Pro-Leu-Leu-Pro-His-Ser-Asn-Leu-Asp-His-Ile-Leu-Glu-Pro-Ser.

7. The kit of claim 4, for confirming the presence
20 of such HTLV-1 infection, wherein said solid support includes a strip having said gp46 peptide antigen immobilized at one position on the strip, and one or more viral antigens effective to positively confirm all HTLV-I and HTLV-II positive sera, immobilized at one or more
25 positions on the strip spaced apart from the position of the gp46 antigen.

8. The kit of claim 4, which further includes an
HTLV-II specific antigen capable of reacting specifically
30 with antibodies from HTLV-II positive sera.

50

9. The kit of claim 8, wherein the HTLV-II specific peptide has an HTLV-II specific immunogenic region formed by the amino acid sequence Met-Thr-Leu-Leu-Val-Asp-Ala-Pro-Gly-Tyr-Asp-Pro-Leu-Trp-Phe-Ile-Thr-Ser-Glu-Pro-Thr-
5 Gln-Pro-Pro-Pro-Thr-Ser-Pro-Pro-Leu-Val-His-Asp-Ser-Asp-Leu-Glu-His-Val-Leu-Thr-Pro-Ser-Thr-Ser-Trp-Thr-Thr-Lys.

10. The kit of claim 9, wherein the HTLV-II specific peptide immunogenic region is formed by the
10 amino acid sequence Ser-Pro-Pro-Leu-Val-His-Asp-Ser-Asp-Leu-Glu-His-Val-Leu-Thr-Pro-Ser-Thr-Ser-Trp-Thr-Thr-Lys.

11. The kit of claim 9, wherein the HTLV-II specific peptide immunogenic region is formed by the
15 amino acid sequence Ser-Pro-Pro-Leu-Val-His-Asp-Ser-Asp-Leu-Glu-His-Val-Leu-Thr-Pro-Ser.

12. A method of detecting HTLV-I positive human sera comprising
20 reacting the sera with a peptide antigen which is immunoreactive with anti-HTLV-I monoclonal antibody derived from ATCC cell line HB8755, and
detecting the presence of human antibody bound to the peptide.

25 13. The method of claim 12, wherein the peptide contains is less than about 77 amino acids derived from HTLV-I envelope protein gp46, and includes the immunogenic region formed by the amino acid sequence Ser-
30 Leu-Leu-Val-Asp-Ala-Pro-Gly-Tyr-Asp-Pro-Ile-Trp-Phe-Leu-Asn-Thr-Glu-Pro-Ser-Gln-Leu-Pro-Pro-Thr-Ala-Pro-Pro-Leu-

51

Leu-Pro-His-Ser-Asn-Leu-Asp-His-Ile-Leu-Glu-Pro-Ser-Ile-Pro-Trp-Lys-Ser-Lys.

14. A method of producing a peptide for use in
5 detecting HTLV-I positive human sera comprising
- (i) generating a mixture of random-sequence polynucleotides of the form: $(N_1N_2N_3)_m$, where $N_1N_2N_3$ represent different trinucleotide codons corresponding to substantially each natural L-amino acid, and m is 5-10,
 - 10 (ii) forming a library of random-sequence vectors by inserting the polynucleotides into a vector capable of expressing the inserted sequence;
 - (iii) manipulating the library vectors to express the random-sequence polynucleotides as random amino acid
15 sequences,
 - (iv) screening the library vectors for the presence of an amino acid sequence which is immunoreactive with anti-HTLV-I monoclonal antibody derived from ATCC cell line HB8755,
 - 20 (v) isolating a library vector which expresses such an immunoreactive amino acid sequence; and
 - (vi) producing the polypeptide encoded by the inserted sequence in the isolated vector.
- 25 15. A peptide antigen comprising less than about 50 amino acids derived from HTLV-II envelope protein gp46, said antigen including the immunogenic region formed by the amino acid sequence Met-Thr-Leu-Leu-Val-Asp-Ala-Pro-Gly-Tyr-Asp-Pro-Leu-Trp-Phe-Ile-Thr-Ser-Glu-Pro-Thr-Gln-Pro-Pro-Pro-Thr-Ser-Pro-Pro-Leu-Val-His-Asp-Ser-Asp-Leu-
30 Glu-His-Val-Leu-Thr-Pro-Ser-Thr-Ser-Trp-Thr-Thr-Lys.

16. The peptide of claim 15, wherein the HTLV-II specific peptide immunogenic region is formed by the amino acid sequence Ser-Pro-Pro-Leu-Val-His-Asp-Ser-Asp-Leu-Glu-His-Val-Leu-Thr-Pro-Ser-Thr-Ser-Trp-Thr-Thr-Lys.

5

17. The peptide of claim 15, wherein the HTLV-II specific peptide immunogenic region is formed by the amino acid sequence Ser-Pro-Pro-Leu-Val-His-Asp-Ser-Asp-Leu-Glu-His-Val-Leu-Thr-Pro-Ser.

10

18. A kit for detecting the presence of HTLV-II infection in a human serum, comprising

a solid support with an immobilized peptide antigen comprising less than about 50 amino acids derived from HTLV-II envelope protein gp46, said antigen including the immunogenic region formed by the amino acid sequence Met-Thr-Leu-Leu-Val-Asp-Ala-Pro-Gly-Tyr-Asp-Pro-Leu-Trp-Phe-Ile-Thr-Ser-Glu-Pro-Thr-Gln-Pro-Pro-Pro-Thr-Ser-Pro-Pro-Leu-Val-His-Asp-Ser-Asp-Leu-Glu-His-Val-Leu-Thr-Pro-Ser-Thr-Ser-Trp-Thr-Thr-Lys, and

20

reporter means for detecting the presence of human antibodies bound to said peptide.

19. A composition for use as a vaccine against infection by HTLV-I comprising

25

a peptide antigen comprising less than about 77 amino acids derived from HTLV-I envelope protein gp46, said antigen including the immunogenic region formed by the amino acid sequence Ser-Leu-Leu-Val-Asp-Ala-Pro-Gly-Tyr-Asp-Pro-Ile-Trp-Phe-Leu-Asn-Thr-Glu-Pro-Ser-Gln-Leu-Pro-Pro-Thr-Ala-Pro-Pro-Leu-Leu-Pro-His-Ser-Asn-Leu-Asp-His-Ile-Leu-Glu-Pro-Ser which is immunoreactive with

30


anti-HTLV-I monoclonal antibody derived from ATCC cell line HB8755; and

(B) a carrier to which the peptide is attached.

- 5 20. The vaccine of claim 19, for use also as a vaccine against infection by HTLV-II, which further includes, attached to the immunogenic carrier, a peptide antigen comprising less than about 50 amino acids derived from HTLV-II envelope protein gp46, said antigen.
- 10 including the immunogenic region formed by the amino acid sequence Met-Thr-Leu-Leu-Val-Asp-Ala-Pro-Gly-Tyr-Asp-Pro-Leu-Trp-Phe-Ile-Thr-Ser-Glu-Pro-Thr-Gln-Pro-Pro-Pro-Thr-Ser-Pro-Pro-Leu-Val-His-Asp-Ser-Asp-Leu-Glu-His-Val-Leu-Thr-Pro-Ser-Thr-Ser-Trp-Thr-Thr-Lys.

1/8

Fig. 1A



5565 MTA4 5790

5664 5807 MTA1

5664 MTA5 5895

Fig. 1B

<p> HILEPSIPWKSLLTLVQL HSNLDHILEPS TAPLLPHSNLDHILEPS </p>	<p> K164 K162 K163 </p>
<p> gt-SLLVDAPGYDPIWFLNTEPSQLPPTAPPLLPHSNLDHILEPSIPWKSLLTLVQLTLQSTJYYCIVCIDRASLSTWHV-gt gt-SLLVDAPGYDPIWFLNTEPSQLPPTAPPLLPHSNLDHILEPSIPWKS-gt CGFPSSLLVDAPGYDPIWFLNTEPSQLPPTAPPLLPHSNLDHILEPS-gt </p>	<p> MTA-5 MTA-1 MTA-4 </p>
<p> CGFPFSLLDAPGYDPIWFLNTEPSQLPPTAPPLLPHSNLDHILEPSIPWKSLLTLVQLTLQSTNYYCIVCIDRASLSTWHVLY- CGSSMTLLVDAPGYDPLWFLTSEPTQPPPTSPPLVHDSBLEHVLTPSTSWTKILKFIQLTLQSTNYYSCMVCVDRSSLSWHVLY- </p>	<p> HTLV-1 HTLV-1I </p>
<p> gt-MTLLVDAPGYDPLWFLTSEPTQPPPTSPPLVHDSBLEHVLTPSTSWTK-gt </p>	<p> GHZ-K15 </p>
<p> EHVLTPTSTSWTT SPPLVHDSBLEHVLTPSTSWTT SPPLVHDSBLEHVLTPS WFLTSEPTQPPPTS MTLLVDAPGYDPLW </p>	<p> K169 K170 K125 K126 K128 </p>

Fig. 3

4/8

Fig. 4A





Fig. 4B



5/8

SfiI SfiI

5' - CTATTCTCACTCGGCCGACG | GGGCCGAAACTGTTGAA - 3'

3' - GATAAGAGTGAGCCGGC | GACCCCGGCTTTGACAACTT - 5'





Fig. 5

Degenerate BglI fragments:

GGGCT (NNK)₇, GGGGCCGCTG
TGCCCCGA (NNM)₇, CCCCAGC




Multiple copies of phage having
recombinant pIII surface protein:

Recombinant pIII:

NH₂ -A D G A X₇ G A A G A E T V E - . . .

6/8

**Fig. 5 (con't)**

7/8

5521 5531 5541 5551 5561 5571
 TGGACATCCGCATACACGGGCCCCGTCTCCAGTCCATCCTGGAAGTTTCATTCAGA

5581 5591 5601 5611 5621 5631
 TGTAAATTTACCCAGGAAGTCAGCCAAGTGTCCCTTCGACTACACTTCTCTAAGTGCGG

M T L L V D A P G Y D P L W F I T S

5641 5651 5661 5671 5681 5691
 CTCCTCCATGACCCTCCTAGTAGATGCCCCTGGATATGATCCTTTATGGTTCATCACCTC
 (KAE1=) ATGACCCTCCTAGTAGATGCC-->

E P T Q P P P T S P P L V H D S D L E H
 5701 5711 5721 5731 5741 5751
 AGAACCCACTCAGCCTCCACCAACTTCTCCCCATTGGTCCATGACTCCGACCTTGAACA

V L T P S T S W T T K

5761 5771 5781 5791 5801 5811
 TGTCCTAACCCCTCCACGTCCTGGACGACCAAAATACTCAAATTTATCCAGCTGACCTT
 <--AGGTGCGGACCACGTGGTTTT =(KAE5)
 CAG^CTG (PvuII)

5821 5831 5841 5851 5861 5871
 ACAGAGCACCAATTACTCCTGCATGGTTTGCGTGGATAGATCCAGCCTCTCATCCTGGCA

5881 5891 5901 5911 5921 5931
 TGTACTCTACACCCCCAACATCTCCATTCCCCAACAAACCTCCTCCCGAACCATCCTCTT

Fig. 6





Fig. 7

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 92/00823

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 C12N15/00; G01N33/566;	C07K7/10; G01N33/577	A61K39/12; A61K39/42
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	C07K ; C12N	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category *	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X	WO,A,8 906 543 (GENELABS INCORPORATED) 27 July 1989	1,4,12, 13,19
Y	see claims 1-10; figures FIG.1A,1B ---	14
X	WO,A,9 008 162 (UNITED BIOMEDICAL INC.) 26 July 1990 see claims ---	3,6
Y	SCIENCE, vol. 249, 27 July 1990, AAAS, WASHINGTON, DC, US; pages 386 - 390; J.K. SCOTT AND G. SMITH: 'Searching for peptide ligands with an epitope library' cited in the application --- -/--	14
<p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
17 JUNE 1992	22. 06. 92	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	HORNIG H. 	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
X	WO,A,9 010 231 (REPLICO MEDICAL AB) 7 September 1990 see claims ---	1-6, 8-13, 15-17
P,X	EP,A,0 439 077 (UNITED BIOMEDICAL INC.) 31 July 1991 see claims see figure 2 ---	1-13, 15-17

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO. US 9200823
SA 57776**

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 17/06/92

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
WO-A-8906543	27-07-89	AU-B-	613350	01-08-91
		AU-A-	1367088	11-08-89
		EP-A-	0395634	07-11-90
		JP-T-	3501722	18-04-91

WO-A-9008162	26-07-90	CA-A-	2025019	14-07-90
		EP-A-	0404935	02-01-91
		JP-T-	3504165	12-09-91

WO-A-9010231	07-09-90	AU-A-	5283490	26-09-90
		EP-A-	0461193	18-12-91

EP-A-0439077	31-07-91	None		
